

Synonym

Spike,S protein RBD,Spike glycoprotein Receptor-binding domain,S glycoprotein RBD,Spike protein RBD

Source

SARS-CoV-2 Spike RBD, His Tag (BA.3/Omicron) (SPD-C522i) is expressed from human 293 cells (HEK293). It contains AA Arg 319 - Lys 537 (Accession # [QHD43416.1](#) (G339D,S371F,S373P,S375F,D405N,K417N,N440K,G446S,S477N,T478K,E484A,Q493R,Q498R,N501Y,Y505H)). The spike mutations are identified on the SARS-CoV-2 Omicron variant (Pango lineage: BA.3; GISAID clade: GRA; Nextstrain clade: 21M).  
Predicted N-terminus: Arg 319

Molecular Characterization

This protein carries a polyhistidine tag at the C-terminus.  
The protein has a calculated MW of 26.8 kDa. The protein migrates as 33-37 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.  
>95% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.  
Contact us for customized product form or formulation.

Reconstitution


Please see Certificate of Analysis for specific instructions.  
*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.  
*Please avoid repeated freeze-thaw cycles.*  
This product is stable after storage at:


- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



SARS-CoV-2 Spike RBD, His Tag (BA.3/Omicron) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.


SEC-MALS




The purity of SARS-CoV-2 Spike RBD, His Tag (BA.3/Omicron) (Cat. No. SPD-C522i) is more than 95% and the molecular weight of this protein is around 38-56 kDa verified by SEC-MALS.  
[Report](#)

Bioactivity-ELISA







Immobilized SARS-CoV-2 Spike RBD, His Tag (BA.3/Omicron) (Cat. No. SPD-C522i) at 2 µg/mL (100 µL/well) can bind Human ACE2, Fc Tag (Cat. No. AC2-H5257) with a linear range of 1-31 ng/mL (QC tested).



Immobilized SARS-CoV-2 Spike RBD, His Tag (BA.3/Omicron) (Cat. No. SPD-C522i) at 1 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (Cat. No. S1N-M122) with a linear range of 0.2-6 ng/mL (Routinely tested).



Immobilized SARS-CoV-2 Spike RBD, His Tag (BA.3/Omicron) (Cat. No. SPD-C522i) at 1 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 Spike RBD Broadly Neutralizing Antibody, Human IgG1 (Cat. No. SPD-M265) with a linear range of 0.001-0.05 µg/mL (Routinely tested).



Immobilized SARS-CoV-2 Spike RBD, His Tag (BA.3/Omicron) (Cat. No. SPD-C522i) at 1 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 Spike RBD Antibody, Mouse IgG1 (Omicron Specific) (Cat. No. SPD-M305) with a linear range of 0.1-25 ng/mL (Routinely tested).

Background

It's been reported that Coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

