




Synonym

Spike,S protein RBD,Spike glycoprotein Receptor-binding domain,S glycoprotein RBD,Spike protein RBD

Source

SARS-CoV-2 S protein RBD (N501Y), His Tag (SPD-C52Hn) is expressed from human 293 cells (HEK293). It contains AA Arg 319 - Lys 537 (Accession # [QHD43416.1](#) (N501Y)). The N501Y mutaion was identified in multiple SARS-CoV-2 Variants of Concerns (VOCs), i.e. the Alpha variant (Pango lineage: B.1.1.7; other names: 20I/501Y.V1 or VOC 202012/01), the Beta variant (Pango lineage: B.1.351; other names: 20H/501Y.V2), the Gamma variant (Pango lineage: P.1; other names: 20J/501Y.V3).  
Predicted N-terminus: Arg 319

Molecular Characterization



This protein carries a polyhistidine tag at the C-terminus.  
The protein has a calculated MW of 26.6 kDa. The protein migrates as 33-35 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.  
>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.  
Contact us for customized product form or formulation.

Reconstitution


Please see Certificate of Analysis for specific instructions.  
*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.  
*Please avoid repeated freeze-thaw cycles.*  
This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



SARS-CoV-2 S protein RBD (N501Y), His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

SEC-MALS




The purity of SARS-CoV-2 S protein RBD (N501Y), His Tag (Cat. No. SPD-C52Hn) is more than 90% and the molecular weight of this protein is around 28-38 kDa verified by SEC-MALS.

[Report](#)




SARS-CoV-2 (COVID-19) S protein RBD (N501Y), His Tag (MALS verified)


Catalog # SPD-C52Hn



Immobilized Human ACE2, Fc Tag (Cat. No. AC2-H5257) at 1 µg/mL (100 µL/well) can bind SARS-CoV-2 S protein RBD (N501Y), His Tag (Cat. No. SPD-C52Hn) with a linear range of 2-39 ng/mL (QC tested).




Immobilized SARS-CoV-2 S protein RBD (N501Y), His Tag (Cat. No. SPD-C52Hn) at 1 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 RBD Potent Neutralizing Antibody, Human IgG1 (Cat. No. SPD-M180) with a linear range of 0.1-3 ng/mL (Routinely tested).



Serial dilutions of Monoclonal Anti-SARS-CoV-2 RBD Potent Neutralizing Antibody, Human IgG1 (Cat. No. SPD-M180) were added into SARS-CoV-2 S protein RBD (N501Y), His Tag (Cat. No. SPD-C52Hn) : Biotinylated Human ACE2, His,Avitag (Cat. No. AC2-H82E6) binding reactions. The half maximal inhibitory concentration (IC50) is 1.13169 (Routinely tested).

Bioactivity-SPR



Human ACE2, Fc Tag (Cat. No. AC2-H5257) captured on CM5 chip via Anti-human IgG Fc antibodies surface can bind SARS-CoV-2 S protein RBD



SARS-CoV-2 (COVID-19) S protein RBD (N501Y), His Tag (MALS verified)

Catalog # SPD-C52Hn



(N501Y), His Tag (Cat. No. SPD-C52Hn) with an affinity constant of 1.24 nM as determined in a SPR assay (Biacore 8K) (Routinely tested).

Background

It's been reported that Coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

Discounts, Gifts,  
and more!

